

COVID-19 Vaccination Schedules: mRNA vaccines

Recipient Age	Product	Persons who ARE NOT Moderately or Severely Immunocompromised ⁺		Persons who ARE Moderately or Severely Immunocompromised ⁺	
		Primary Series	Booster Dose(s)	Primary Series	Booster Dose(s)
5-11 years	Pfizer	2 doses, at least 3 weeks apart	Not recommended	3 doses	Not recommended
12-17 years	Pfizer	2 doses, 3-8 weeks apart	At least 5 months after primary series	3 doses	1 st : At least 12 weeks after primary series 2 nd : At least 4 months after 1 st booster
18-49 years	P/M	2 doses, 3-8 weeks apart (P), 4-8 weeks apart (M)	At least 5 months after primary series	3 doses	1 st : At least 12 weeks after primary series 2 nd : At least 4 months after 1 st booster
50 years+	P/M	2 doses, 3-8 weeks apart (P), 4-8 weeks apart (M)	1 st : At least 5 months after primary series 2 nd : At least 4 months after 1 st booster	3 doses	1 st : At least 12 weeks after primary series 2 nd : At least 4 months after 1 st booster

⁺<https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/immuno.html#mod>

COVID-19 Vaccination Schedules: J&J

Recipient Age	Product	Persons who ARE NOT Moderately or Severely Immunocompromised [†]		Persons who ARE Moderately or Severely Immunocompromised [†]	
		Primary Series	Booster Dose(s)	Primary Series	Booster Dose(s)
18+ years	J&J	1 dose	#1: At least 8 weeks after primary series* #2: At least 4 months after 1 st booster* *mRNA boosters recommended	2 doses, dose #2 should be an mRNA vaccine	At least 8 weeks after primary series* *mRNA boosters recommended

[†]<https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/immuno.html#mod>